

## Screening of the potentiality of four green microalgae to be used as feedstock for biodiesel and nutraceutical production

Soha Mohammed, Hamed Eladel\*, Mohamed Battah and Yasmine

Ibrahim

*Botany and Microbiology Department, Faculty of Science, Benha University,*

*Benha 13518, Egypt.*

### **Abstract:**

Detection of oleaginous microalgae with high lipid productivity is a crucial feature for biofuel and nutraceutical production. Therefore, screening of microalgae from different habitats has become highly significant to face the request on these bioresources. Accordingly, the present study aimed to screen the efficiency of biodiesel feedstock and the ability to accumulate essential fatty acids of four green microalgae (*Chlorobion braunii*, *Tetrademus obliquus*, *Monoraphidium minutum*, and *Asterarcys quadricellulare*), isolated from soil samples of Benha, Egypt. Although *T.obliquus* was the highest biomass productive species ( $114.86 \text{ mgL}^{-1}\text{d}^{-1}$ ), but *M. minutum* was recorded as the highest species of lipid productivity ( $29 \text{ mg L}^{-1}\text{d}^{-1}$ ). Fatty acid profiles of the four microalgae confirmed their suitability for biodiesel production as mentioned by the biodiesel properties within the limits of international standards. Species selection through PROMETHEE and GAIA analysis revealed that *C. braunii* is the best promising as a potential source for biodiesel production because it has a high saturated fatty acids proportion (39.75%) and shows cetane number (54.99) and low iodine value (100.57) and many other biodiesel properties within the standards limit. In addition, the screened microalgae showed a high percentage of omega fractions in their total fatty acids, whereas *Asterarcys quadricellulare* and *M. minutum* have high omega-3 fatty acids content of 34.4% and 28.9%, respectively.

**Keywords:** Green microalgae, Biodiesel, Lipid, Essential Fatty Acids, Nutraceuticals.

## ***Introduction***

Microalgae include a varied group of photoautotrophic microorganisms that inhabit aquatic habitats and terrestrial ecosystems. A long time ago, the screening of microalgae attracted attention in respect of their high potency for biotechnological applications including food supplements to prevent many diseases, and biodiesel production (**Bhalamurugan et al. 2018; Khan et al. 2018**). Microalgae show great potential as feedstocks in replacing fossil fuels to ensure energy security and to avoid environmental pollution globally (**Enamala et al. 2018**). Biodiesel raw materials obtained from microalgae appeared to be one of the foremost promising alternative sources for biodiesel generation due to their rapid growth rates, high biomass productivity and high lipid contents over the terrestrial crop plants. Microalgae produce a significantly higher yield of lipids, about 150 000 L of oil per hectare, which is three times higher than other oil crops (**Li et al. 2020**). In addition, microalgae can grow on wastewaters and adapt themselves to adverse environmental conditions (**Eladel et al. 2019**). Also, with their photosynthetic effectiveness and CO<sub>2</sub> sequestration, they can be grown successfully with high lipid content appropriate as biodiesel feedstock and do not compete with human nutrition (**Almarashi et al. 2020**). A large number of oleaginous microalgae species have already been substituted a fraction of petroleum-based fuels in some countries (**Meng et al. 2009**). Typical microalgae for biodiesel generation require not only high lipid content but also reasonable fatty acids composition. The generation of saturated and monounsaturated fatty acids by microalgae is ideal for biodiesel production, where it achieves the equilibrium between cetane number and cold-flow properties (**Koller et al. 2014; Ma et al. 2014**).

As a result of algal growth in diverse and extreme habitats, they produce a group of unique bioactive compounds. Polyunsaturated fatty acids (PUFAs) are highly beneficial bioactive products for human health. Algae have a useful effect on human nutrition and health, because of their high polyunsaturated fatty acids (**Morais et al. 2015**). Algal lipids are of an extraordinary commercial value as sources of nutritiously important polyunsaturated fatty acids (PUFAs) and are therefore broadly utilized as ingredients in the nutraceutical industry (**Mendis and Kim 2011**). Replacing saturated lipids with unsaturated is effective in preventing cardiovascular and cancer diseases (**Clifton and Keogh 2017**). The human body cannot generate essential fatty acids, and hence, these must be consumed in the diet. They are advisable for the proper growth and development of organisms. Omega-3 fatty acids alpha-linolenic acid (ALA), eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) provide substantial protection against chronic and metabolic diseases, therefore, their intake is advisable (**Bazinet and Layé 2014**). Dietary omega-3 PUFAs enhance the body functions to fight against inflammatory conditions that lead to initiating diseases like cancer, kidney malfunction, diabetes, and cardiovascular disorders (**Yagi et al. 2017**). Docosahexaenoic (DHA) is a vital bioactive lipid that is suggested for keeping up physiological health. In addition, docosahexaenoic acid (DHA) contributes to the upregulation of intracellular antioxidants and driving oxidative stress reduction. Other than this, DHA is additionally reported to display an anti-inflammatory effect (**Aluko 2012**). The valuable omega-6 fatty acids are linoleic acid (LA), gamma-linolenic acid (GLA), arachidonic acid (AA) and docosapentaenoic acid, (DPA) are the major fatty acids of this group. They play a significant part in numerous physiological functions and molecular pathways (**Maki et al. 2018**). Linoleic acid reduces the risk of blood pressure and activates prostaglandins

synthesis. It leads to the composition of arachidonic acid, eicosapentaenoic acid, and docosahexaenoic acid that play principal roles in regulating body homeostasis (**Ward and Singh 2005; Aluko 2012**).

Recently, microalgae biorefinery models have been applied to reduce biodiesel production costs through biomass valorization. For example, lipid-free biomass residue after lipid extraction can be utilized as a nourishing supplement (**Abomohra et al. 2018**). This study intended to isolate some local soil microalgae and evaluate their potential ability for utilizing them in biodiesel production and food supplements. Also, the study aims to determine the most suitable species as biodiesel feedstock by PROMETHEE and GAIA software (**Islam et al. 2013; Kramadibrata et al. 2019**).

## ***Materials and Methods***

### ***Microalgae isolation and growth conditions***

The soil samples have been collected from agricultural land in Benha city (coordinates, 30°25'47.4"N 31°12'55.2"E), Egypt. Ten grams of soil were mixed with 50 mL of sterilized distilled water. The soil suspension was agitated at 300 rpm for 30 min, then 10-fold serial dilutions (**Anderson, 2005**) were prepared using Bold's Basal medium (BBM). Using the spread agar method; Petri-dishes containing solid BBM were inoculated with different soil dilutions and incubated at  $26 \pm 2.0$  °C. Cultures were illuminated by tubular fluorescent lamps at  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$  light intensity. After microalgal growth appeared, the individual colonies were picked up and sub-cultured several times on agar plates until the complete purification of the isolates. Pure microalgal isolates were examined using a light

microscope, re-cultivated in 50 mL liquid BBM, and incubated under the aforementioned growth conditions. Microalgae were cultivated in batch culture containing 700 mL BBM medium. All flasks were autoclaved, cooled down, and inoculated with exponentially grown culture to get an initial optical density ( $OD_{680}$ ) of  $0.05 \pm 0.003$ . All cultures were aerated using an atmospheric air pump. Microalgal identification was conducted morphologically using the classification systems adopted by the following references: **Hegewald & Schmidt (1992)**, **Komárek & Fott (1983)** and **Wehr *et al.* (2015)**.

### ***Growth measurements***

Batch cultures were incubated for up to 14 days, and the microalgae growth was monitored by measuring the optical density ( $OD_{680}$ ) every two days interval (**Held, 2011**). Dry weight was determined at the early exponential and late exponential phases of each microalgal species for biomass productivity determination. To measure the dry weight, algal cells in 10 mL culture were collected by filtration through 0.45  $\mu$ m pre-weighted filter paper, then oven-dried at 105 °C until constant weight. Biomass productivity determination using equation (1);

$$\text{Biomass productivity (mg L}^{-1}\text{d}^{-1}\text{)} = (dwt - dwi)/\Delta t \quad (1)$$

Where *dwt* and *dwi* represent the cellular dry weight at the measured time (*t*) and the initial dry weight, respectively, while  $\Delta t$  represents the time interval in days (**Eladel *et al.* 2019**)

***Lipid content***

Microalgae lipid extraction was carried out according to the modified **Folch et al. (1957)** method. Briefly, 20 mL of microalgal culture were centrifuged at 3000× g for 10 minutes. The supernatant was disposed of, at that point, 10 mL of chloroform/methanol (2/1, v/v) were added to the cell pellet and let at room temperature for two hours on a shaker at 200 rpm. The homogenate was centrifuged at 3000× g for ten minutes to recover the liquid phase, which was transferred to a clean tube. To facilitate the separation of water-soluble components, 2 mL of 0.9% NaCl were added and vortexed vigorously for 30 sec., then the blend was centrifuged at 200× g for 2 min to partition into two phases. The lower phase was pulled to a pre-weighed vial, the solvent was released, dried at 80 °C for 30 min, and weighed to determine the lipid content. Microalgae lipid productivity was calculated in the exponential phase of growth according to equation (2);

$$\text{Lipid productivity (mg L}^{-1}\text{d}^{-1}) = (LCt - LCi)/\Delta t \quad (2)$$

Where  $LCt$  and  $LCi$  represent lipid production ( $\text{mg L}^{-1}$ ) after the time ( $t$ ) and the initial lipid content, respectively, while  $\Delta t$  represents the time interval in days.

***Fatty acids analysis***

Biodiesel was prepared from the lipid extract according to a modified method by **Christie (1993)**. Lipid extracts were dried beneath a stream of argon,

at that point 333  $\mu\text{L}$  of methanol: toluene (1:1, v/v) and 167  $\mu\text{L}$  of 0.5 M sodium methoxide were added and let for 20 min at room temperature. After incubation, 500  $\mu\text{L}$  of 1 M NaCl and 50  $\mu\text{L}$  of 37% HCl were included in the mixture and vortexed vigorously for 30 sec. To extract FAMES, 1.5 mL of hexane were added and vortexed vigorously. The mixture was centrifuged at  $200\times g$  for 2 min to isolate the two phases. The upper hexane phase was pulled to a 2 mL Eppendorf tube, at that point the solvent was released beneath a stream of argon. FAMES were dissolved in 40  $\mu\text{L}$  of acetonitrile and analyzed by GC-FID equipped with 30 m column (J&W HP-5, 0.32 mm diameter, and 0.25  $\mu\text{m}$  inner film) using helium as a carrier gas. Injector and detector temperatures were kept at 280  $^{\circ}\text{C}$  and 300 $^{\circ}\text{C}$ , respectively. The oven temperature program was started at 140  $^{\circ}\text{C}$  for 3 min, increased at 5  $^{\circ}\text{C min}^{-1}$  until 240  $^{\circ}\text{C}$ , then increased at 4  $^{\circ}\text{C min}^{-1}$  until 260  $^{\circ}\text{C}$  where it was fixed at this temperature for 5 min.

### ***Biodiesel properties based on FAME profiles***

Because of the need for large amounts of biodiesel and specialized instrumentation, which are not available. So, Predictive models based on FA composition were used for the calculation of the screened microalgae biodiesel properties. The main biodiesel quality parameters include an average degree of unsaturation (ADU), kinematic viscosity (KV), specific gravity (SG), cloud point (CP), cetane number (CN), iodine value (IV), and higher heating value (HHV) were determined by mathematical models (**Hoekman et al. 2012; Song et al. 2013**) per equations (3-9). However, cold filter plugging point (CFPP) and long-chain saturation factor (LCSF) were calculated according to **Ramos et al. (2009)**

using equations (10-11). Predictive oxidative stability (OS); was calculated using equation (12) according to **Park et al. (2008)**.

$$ADU = \sum M \times Y_i \quad (3)$$

Where M is the number of carbon-carbon double bonds in FA constituent and  $Y_i$  is the mass fraction of each FA constituent, respectively.

$$\text{Kinematic viscosity} = -0.6313X + 5.2065 \quad (4)$$

$$\text{Specific gravity} = 0.0055X + 0.8726 \quad (5)$$

$$\text{Cloud point} = -13.356X + 19.994 \quad (6)$$

$$\text{Iodine value} = 74.373X + 12.71 \quad (7)$$

$$\text{Cetane number} = -6.6684X + 62.876 \quad (8)$$

$$\text{HHV} = 1.7601X + 38.534 \quad (9)$$

Where X is the ADU.

$$\text{LCSF} = (0.1 \times C16) + (0.5 \times C18) + (1 \times C20) + (1.5 \times C22) + (2 \times C24) \quad (10)$$

$$\text{CFPP} = (3.1417 \times \text{LCSF}) - 16.477 \quad (11)$$

$$\text{OS (h)} = (117.9295/C18p) + 2.5905 \quad (12)$$

C18p represents the content (weight %) of linoleic (C18:2) and linolenic acids (C18:3).

### ***Microalgae selection based on biodiesel properties***

The screened microalgae were ranked based on biodiesel properties to select the most suitable microalgae species for biodiesel production using PROMETHEE and GAIA software v1.4.0.0. Preference Ranking Organization Method for Enrichment of Evaluation (PROMETHEE) and Graphical Analysis for Interactive Aid (GAIA) (Islam et al. 2013, Kramadibrata et al. 2019).


### ***Statistical analysis***

The results were calculated as the mean  $\pm$  standard deviation (SD), where each experiment was performed in three replicates. In addition, SPSS software (IBM, v20) was used to perform a one-way (ANOVA) at a probability level ( $P \leq 0.05$ ).

## ***Results and Discussion***

### ***Microalgal identification***

The screened unialgal isolates were purely isolated through agar plate's technique and sub-culturing in a liquid media. Using microscopic characterization, four green microalgae species were morphologically identified as *Chlorolobion braunii* (formerly *Monoraphidium braunii*), *Tetradesmus obliquus* (formerly *Scenedesmus obliquus*), *Monoraphidium minutum* (formerly *Selenastrum minutum*), and *Asterarcys quadricellulare* (formerly *Coelastrum quadricellulare*) (Fig. 1).




**Fig. 1:** Photomicrographs of the four isolated microalgae, (a) *Chlorolobion braunii*, (b) *Tetradesmus obliquus*, (c) *Asterarcys quadricellulare*, (d) *Monoraphidium minutum*.  
Scale bars: 10  $\mu\text{m}$  (a, b, d); 20  $\mu\text{m}$  (c).

### ***Microalgal growth and lipid content***


Microalgae are extensively discussed as a valuable source for third-generation biofuels, especially as a sustainable and environmentally-friendly feedstock for biodiesel production. In the present study, four species of soil green microalgae were screened based on their lipid productivity. All screened species showed a lag phase of two days before they began their exponential growth. Growth curves of the four screened microalgae revealed their exponential growth phase in between the fourth and tenth day of incubation time (Fig. 2). *Tetradesmus* significantly recorded the highest biomass productivity  $114.86 \text{ mg L}^{-1} \text{ d}^{-1}$ , while *Chlorolobion* was the lowest biomass productive ( $35.14 \text{ mg L}^{-1} \text{ d}^{-1}$ ) compared to the other species (Fig. 3). At the exponential phase of microalgae, lipid productivity revealed that *Monoraphidium* was significantly the highest lipid productive species with up to  $29.1 \text{ mg L}^{-1} \text{ day}^{-1}$ . In contrast, as compared to certain other screened microalgae, *Asterarcys* microalga had significantly lower lipid productivity (Fig. 4).

### ***Microalgae fatty acids profile***


Fatty acids profile of the screened green microalgae characterized by GC and their relative percentages are recorded in Table 1. Generally, the detected 26 fatty acids with carbon chains ranging from (C11-C24) and different degrees of unsaturation. The most prevailing fatty acids were palmitic acid (17-26%), palmitolic acid (7-12%), oleic acid (7-16%), linoleic acid (4-17%) and alpha-linolenic acid (7-24%), which set within C16 and C18 fatty acids.



**Fig. 2:** Growth curves based on (a) optical density (OD<sub>680nm</sub>) and (b) chlorophyll a, of the four screened microalgae grown in liquid Bold's Basal medium for 14 days. Error bars show the standard deviation of three replicates



**Fig. 3:** Biomass productivity of the four screened microalgae at exponential phase. Error bars show the SD for three replicates. Columns of the same series with the same letter showed insignificant difference at  $P \leq 0.05$



**Fig. 4:** Lipid content and Lipid productivity of the four screened microalgae at the exponential phase. Error bars show the SD for the three replicates. The same series with the same letter showed insignificant difference at  $P \leq 0.05$

Palmitic acid (C16:0) and stearic acid (C18:0) are two of the most commonly observed fatty acids synthesized by microalgae and are suitable for biodiesel production (**Díaz et al. 2014**). The degree of unsaturation plays a critical role in biodiesel properties as the higher the degree of unsaturation of the FAMES, the higher the oxidation stability of the biodiesel (**Islam et al. 2013**). The present results revealed that *A. quadricellulare* recorded the highest percentage of unsaturation fatty acids (75.57%), while *C. braunii* recorded the lowest value (60.27%). In addition, all of our isolates were characterized by a high percentage of PUFAs more than MUFAs (Table 1).

### ***Microalgal biodiesel properties***

Biodiesel properties analysis of a biodiesel sample requires a long time and a lot of money. In some cases, it may be impossible to obtain a sufficiently large sample of biodiesel from an arising feedstock oil for detailed analyses, such as algal biodiesel. Mathematical models were applied to predict the properties of biodiesel depending on the fatty acid profile. Subsequently, it may be utilized as a bioprospecting tool for rapidly estimating the potential utility of a new feedstock (**Talebi et al. 2013**). The most important properties of biodiesel as a substitute for diesel fuel are the cetane number, viscosity, and density, cold filter plugging point, oxidative stability, ignition quality, combustion heat and cold flow (**Ashour et al. 2019**). Because the calculated values using the equations are closer to the practical analyzed biodiesel values (**Song et al. 2013, Ma et al. 2014**). The biodiesel characteristics of the screened microalgae were compared to the international standards, where it showed ideal cetane number, viscosity, specific gravity, and oxidation stability (Table 2).

**Table 1: Fatty acids composition of the four screened green microalgae species (% of total fatty acids)**

Fatty acid	<i>Chlorobion braunii</i>	<i>Tetrademus obliquus</i>	<i>Monoraphidium minutum</i>	<i>Asterarcys quadricellulare</i>
C11:0 Undecanoic acid	0.69	0.21	0.3	0.34
C13:0 Tridecanoic acid	1.42	1.37	0.9	1.17
C14:0 Myristic acid	1.21	1.23	1.19	0.87
C15:0 Pentadecanoic acid	2.67	3.88	2.4	2.94
C16:0 Palmitic acid	25.98	23.71	16.75	17.17
C17:0 Heptadecanoic acid	5.26	2.39	4.73	0.95
C18:0 Stearic acid	1.31	1.37	1.05	1.0
C24:0 Lignoceric acid	1.21	0.48	0.7	ND
<b>Total saturated fatty acids (SAF)</b>	<b>39.75</b>	<b>34.64</b>	<b>28.02</b>	<b>24.44</b>
C16:1 ω9 Palmitolic acid	10.28	11.45	7.04	8.97
C16:1 ω7 Palmitoleic acid	1.99	2.12	1.72	3.55
C18:1 ω9 Oleic acid	12.31	7.21	13.52	16.43
C18:1 ω7 Vaccenic acid	1.29	1.39	1.92	2.43
C20:1 ω9 Gondoic acid	0.34	ND	ND	0.23
C24:1 ω9 Nervonic acid	2.15	1.59	0.7	0.89

<b>Total monounsaturated fatty acids (MUFA)</b>	<b>28.36</b>	<b>23.76</b>	<b>24.9</b>	<b>32.5</b>
C16:2 ω4 Hexadecadienoic acid	ND	ND	0.29	ND
C16:3 ω4 Hexagonic acid	1.33	1.33	0.96	1.23
C16:4 ω3 Hexadecatetraenoic acid	2.59	6.01	10.4	10.06
C18:2 ω6 Linoleic acid (LA)	16.7	11.67	11.68	3.68
C18:3 ω6 γ-linolenic acid	0.43	0.9	0.45	0.18
C18:3 ω4 Octadecadienoic acid	ND	0.86	ND	0.32
C18:3 ω3 α-Linolenic acid (ALA)	6.5	14.79	18.34	24.32
C18:4 ω1 Octadecatetraenoic acid	1.11	2.45	3.78	2.87
C20:4 ω6 Arachidonic acid (AA)	ND	1.38	ND	0.41
C20:5 ω3 Eicosapentaenoic acid (EPA)	0.19	0.22	ND	ND
C22:5 ω6 Docosapentaenoic acid	2.33	1.46	0.3	ND
C22:6 ω3 Docosahexaenoic acid (DHA)	0.73	0.51	0.19	ND
<b>Total polyunsaturated fatty acids (PUFA)</b>	<b>31.91</b>	<b>41.58</b>	<b>46.39</b>	<b>43.07</b>

---

ND = not detected

Results of the present study also revealed that biodiesel characteristics are in agreement with some other related studies (**Islam et al. 2013; Eladel et al. 2019; Abomohra et al. 2022**).

The present study recorded that the content of saturated fatty acids ranged from 24 - 40% (Table 1). It was reported that the more saturated fatty acids supply biodiesel with a higher cetane number, high oxidative stability, decreased emission of nitrous oxide, and improved ignition (**Antolin et al. 2002**). High cetane number led to better combustion, low emission of nitrous oxide and easier engine start-up (**Mandotra et al. 2014**). Cetane numbers for screened microalgae ranged from 51.45 to 54.99, which falls within the limits of the international standards.

High iodine value is always accomplished with a high unsaturation degree and high susceptibility to oxidation and it can be improved by antioxidants addition. While low iodine value offers appropriate oxidation stability (**Islam et al. 2013**). The present results recorded that *Chlorolobion* species has the lowest iodine value and high oxidative stability in comparison with the other screened microalgae. Specific gravity and viscosity are important parameters because they affect the efficiency of fuel atomization. Specific gravity results recorded 0.88 while kinematic viscosity values (4.12 - 4.46) comply with the international standards. One of the main troubles associated with the use of biodiesel is the bad flow properties in cold weather. The present study recorded a relatively high PUFAs content, especially linoleic acid (3.7 - 16.7%), and lower CFPP (-16.4), which agree with standards limits and are in favor of colder environments. Within the present research, the levels of stearic acid C 18:0 (Table 1) were generally very low (below 1.37%). These low values of stearic acid may have contributed to the lower temperatures of CFPP for the studied species (**Tiwari et al. 2021**). The

long-chain PUFAs, especially linoleic acid, resulted in amelioration for biodiesel liquefaction value like CFPP which is in demand for colder environments (Folayan *et al.* 2019; Abomohra *et al.* 2022). In agreement with Santhakumaran *et al.* (2019), a comparison for the biodiesel properties of the four screened microalgae species (Table 2) revealed that *Chlorolobion* lipid extract was the most suitable for biodiesel production. Where it has the highest characteristic biodiesel properties such as the ratio of both saturated fatty acids and monounsaturated fatty acids to the polyunsaturated fatty acids (2.13), the highest CN value (54.99), the highest score in LCSF (0.06%), and the lowest IV value (100.57).

#### ***Selection of suitable microalgae for biodiesel***

Lipid content may vary from organism to organism, where a particular species may have high biomass with a low lipid content and vice versa. So, screening and selection of suitable microalgal species based on various biodiesel properties require a multi-criterion decision analysis (MCDA). (PROMETHEE) and (GAIA) have significant advantages compared to other MCDA methods because it is a promising method, i.e. the decision vector stretch toward the preferred solution (Brans and Mareschal, 2005). In this study, based on phi value, the calculated outranking flows of the tested microalgae pointed out that the most appropriate microalgae for biodiesel production in descending order are *Chlorolobion braunii*, *Tetradasmus obliquus*, *Monoraphidium minutum*, *Asterarcys quadricellulare* (Table 3). GAIA plane is a descriptive supplement to the PROMETHEE rankings.


**Table 2: Comparison of biodiesel properties for lipids of the four screened microalgae and American Society for Testing and Materials (ASTM, 2008), European standards (CEN, 2008)**

	Microalgae of this study				Biodiesel standards	
	<i>Chlorolobion</i>	<i>Tetradesmus</i>	<i>Monoraphidium</i>	<i>Asterarcys</i>	ASTM D6751-08	CEN 14214
Unsaturation degree	1.18	1.51	1.67	1.71	-	-
Kinematic viscosity (mm <sup>2</sup> s <sup>-1</sup> )	4.46	4.25	4.15	4.12	1.9-6.0	3.5-5.0
Specific gravity (kg <sup>-1</sup> cm <sup>3</sup> )	0.88	0.88	0.88	0.88	0.88	0.86-0.9
Cloud point (°C)	4.22	-0.22	-2.37	-2.89	-3 to 12	-
Iodine value (g I <sub>2</sub> 100g <sup>-1</sup> )	100.57	125.27	137.25	140.16	-	Max. 120
Cetane number	54.99	52.78	51.71	51.45	Min. 47	51-120
HHV (MJkg <sup>-1</sup> )	40.61	41.20	41.48	41.55	-	42
Oxidation stability (h)	7.58	6.77	6.46	6.73	Min. 3	Min. 6
LCSF (wt%)	0.06	0.04	0.02	0.02	-	-
CFPP (°C)	-16.30	-16.35	-16.41	-16.41	-13 to -5	-20 to 5

**Table 3: PROMETHEE ranking the four screened green microalgae species based on the Phi scores**

Rank	Microalgae	Phi
1	<i>Chlorolobion braunii</i>	0.0466
2	<i>Tetradesmus obliquus</i>	-0.0020
3	<i>Monoraphidium minutum</i>	-0.0211
4	<i>Asterarcys quadricellulare</i>	-0.0235

The blue lines and diamonds represent the biodiesel quality parameters, the green circles represent the microalgae samples analyzed, i.e. actions; and the red line represents the decision vector. The species along with the decision axis, i.e. *Chlorolobion braunii* microalga is the most suitable candidate among the other species (Fig. 5). Most of the biodiesel properties lie adjacent to the decision axis (Du, KV, SG, CP, CN, HHV, LCSF, and CFPP), while some lie opposite to the decision axis and have less influence on the decision axis.



**Fig. 5:** Graphical Analysis for Interactive Assistance (GAIA) plot of the four screened green microalgae species from the present study showing decision vector

### ***Omega fatty acids from microalgae***

Microalgae have been demonstrated as a prospective source of value-added bioactive compounds with the high commercial value used in pharmaceutical, healthcare and food industries (**Mathimani and Pugazhendhi 2019**). The nutritional value of microalgae is highly related to their essential fatty acid contents (**Liang et al. 2006**). Among value-added products of microalgae, long-chain PUFAs are broadly known for their useful impacts on human health (**Guihéneuf and Stengel 2013**), as they have an important biological role in maintaining a healthy brain function and coronary artery (**Hishikawa et al. 2017**).

Fatty acids of the screened microalgae revealed that oleic acid has the greatest values within the group of omega-9 fatty acids, which ranged from 7.2% to 16.4% (Table 1). Oleic acid is associated with the prevention of brain-related disorders (**Chackalamannil et al. 2017**). Palmitoleic acid an omega-7 MUFA fatty acid which ranged from 1.7% to 3.6% plays a significant role in human metabolism. Researchers have correlated dietary palmitoleic acid with decreased risks of cardiovascular diseases, diabetes, and inflammation (**Wu et al. 2020**). Results presented in Table 4 elucidated that fatty acids profiles of the screened microalgae are characterized by a high level of omega-9 fatty acids (20.3-26.3%), while omega-7 fatty acids percentage (3.3-6%), which does not exceed 1.9 % in many edible plant sources. MUFA fatty acid utilization is appeared to lower blood glucose in individuals with type II diabetes (**Chen et al. 2013**).

PUFAs are divided into two major groups i.e., Omega-3 and omega-6, which play a principal role in the formation of the structure and function of nervous and visual systems of humans and have a protective impact against inflammatory and cardiovascular diseases (**Wysoczański et al. 2016**). Interestingly, most of our microalgae contained a high percentage of omega-3 fatty acids, which exceeds that of many oil sources (Table 4). For example, their percentage was in *A. quadricellulare* recorded 34.38 %, while it was 25.14% in fish oil (**Babalola and Apta 2011**).

In general, omega-3 fatty acids represent the largest proportion of unsaturated fatty acids in most isolates, especially *A. quadricellulare* and *M. minutum*. Moreover, the predominant active omega-3 fatty acid in most isolates is  $\alpha$ -linolenic acid, which constitutes about a quarter of fatty acids content in *A. quadricellulare* compared to other sources of lipids, similar results in which the soil algae *Protosiphon botryoides* produced fatty acids with high PUFAs and high omega-3 fatty acids (**Abomohra et al. 2022**).

Microalgae are the richest source of PUFAs particularly, EPA and DHA, which are the foremost vital PUFA's due to their metabolic functions (**Yen et al. 2013**). DHA and EPA are synthesized by microalgae under various culture conditions (**Adarme-Vega et al. 2012**). **Fraeye et al. (2012)** concluded that feed supplementation with omega-3 PUFAs within the form of fish oil or microalgae is more useful compared to supplementation with ALA through flaxseed oil due to the generally low transformation of ALA to EPA and DHA.

**Table 4: Omega fatty acids percentage of the four tested microalgae in comparison with some other lipid sources**

lipid source	$\omega$ -3	$\omega$ -6	$\omega$ -7	$\omega$ -9	Total	References
<i>Asterarcys quadricellulare</i>	34.4	4.3	6.0	26.3	70.9	Present study
<i>Monoraphidium minutum</i>	28.9	12.1	3.6	21.3	66.0	
<i>Tetradesmus obliquus</i>	21.5	15.4	3.5	20.3	60.7	
<i>Chlorobion braunii</i>	10.0	19.46	3.3	25.1	57.8	
Fish oil	25.1	19.4	10.1	23.3	77.8	Babalola and Apata, (2011)
Palm oil	0.3	11.2	0.3	43.4	55.2	Dorni et al. (2018)
Soybean oil	-	6.1	-	46.6	52.6	Babalola and Apata, (2011)
Ghee (Clarified butter)	0.6	2.0	1.9	23.2	27.6	Dorni et al. (2018)
Coconut oil	0.3	10.5	0.3	4.3	15.2	Babalola and Apata, (2011)

### Conclusion

The present study evaluated the four locally isolated green microalgae based on their fatty acid profiles for biodiesel and nutraceutical production. Generally, the results confirmed that the lipids of these microalgae are suitable as biodiesel feedstock to be performed in a cold climate. Also, it pointed out that *Chlorobion braunii* microalga is the best suitable candidate among the three

other species. The results concluded that the lipids of these microalgae are a convenient nutraceutical source because of their high percentage of essential PUFAs (omega fatty acids), which could be applied in the nutraceutical industry. In addition, because the cost of producing compounds from microalgae is expensive, it is desirable to develop a low-cost bioreactor system to be profitable. Also, growth optimization for maximum biomass and metabolite output is necessary for large-scale applications. Furthermore, the use of wastewater for microalgal biomass production is another area that needs improving to make biotechnological applications cost-effective.

### ***Acknowledgements***

The authors are grateful to the Botany and Microbiology Department, Faculty of Science, Benha University, Egypt, for supporting and providing the facilities to conduct this research. Special thanks to Dr Abdullah A. Saber, Associate Professor of Phycology at the Botany Department, Faculty of Science, Ain Shams University, for his kind assistance in identifying the microalgal species.

### ***References***

**Abomohra, A. E. F., Eladel, H., and Mohammed, S. (2022).** Dual use of a local *Protosiphon* isolate BENHA2020 for biodiesel production and antioxidant activity of lipid-free biomass: A novel biorefinery approach for biomass valorization. *Renewable Energy*.

- Abomohra, A. E. F., Eladel, H., El-Esawi, M., Wang, S., Wang, Q., He, Z., ... and Hanelt, D. (2018).** Effect of lipid-free microalgal biomass and waste glycerol on growth and lipid production of *Scenedesmus obliquus*: Innovative waste recycling for extraordinary lipid production. *Bioresource technology*, **249**, 992-999.
- Adarme-Vega, T. C., Lim, D. K., Timmins, M., Vernen, F., Li, Y., and Schenk, P. M. (2012).** Microalgal biofactories: a promising approach towards sustainable omega-3 fatty acid production. *Microbial cell factories*, **11**(1), 1-10.
- Almarashi, J. Q., El-Zohary, S. E., Ellabban, M. A., and Abomohra, A. E. F. (2020).** Enhancement of lipid production and energy recovery from the green microalga *Asterarcys quadricellulare* by inoculum pretreatment with low-dose cold atmospheric pressure plasma (CAPP). *Energy Conversion and Management*, **204**, 112314.
- Aluko, R. (2012).** Bioactive peptides. In *Functional foods and nutraceuticals* (pp. 37-61). Springer, New York, NY.
- Anderson, R. A. (Ed.). (2005).** *Algal culturing techniques*. Elsevier.
- Antolin, G., Tinaut, F. V., Briceno, Y., Castano, V., Perez, C., and Ramirez, A. I. (2002).** Optimisation of biodiesel production by sunflower oil transesterification. *Bioresource technology*, **83**(2), 111-114.
- Ashour, M., Elshobary, M. E., El-Shenody, R., Kamil, A. W., and Abomohra, A. E. F. (2019).** Evaluation of a native oleaginous marine microalga *Nannochloropsis oceanica* for dual use in biodiesel production and aquaculture feed. *Biomass and Bioenergy*, **120**, 439-447.

- ASTM (2008).** Standard specification for biodiesel fuel blend stock (B100) for middle distillate fuels, ASTM D6751- 08. ASTM International, West Conshohocken
- Babalola, T. O. O., and Apata, D. F. (2011).** Chemical and quality evaluation of some alternative lipid sources for aqua feed production. *Agriculture and Biology Journal of North America*, **2(6)**, 935-943.
- Bazinet, R. P., and Layé, S. (2014).** Polyunsaturated fatty acids and their metabolites in brain function and disease. *Nature Reviews Neuroscience*, **15(12)**, 771-785.
- Bhalarurugan, G. L., Valerie, O., & Mark, L. (2018).** Valuable bioproducts obtained from microalgal biomass and their commercial applications: A review. *Environmental Engineering Research*, **23(3)**, 229–241. <https://doi.org/10.4491/eer.2017.220>
- Brans, J. P., Mareschal, B., (2005).** PROMETHEE methods, multiple criteria decision analysis: state of the art surveys, **163–186**.
- CEN (2008).** (European Committee for Standardization) Automotive fuels: fatty acid methyl esters (FAME) for diesel engines: requirements and test methods, EN14214. European Committee for Standardization, Austrian Standards Institute, Vienna, Austria
- Chackalamannil, S., Rotella, D., and Ward, S. (2017).** *Comprehensive medicinal chemistry III*. Elsevier.
- Chen, B., McClements, D. J., and Decker, E. A. (2013).** Design of foods with bioactive lipids for improved health. *Annual review of food science and technology*, **4**, 35-56.

- Christie, W. W. (1993).** Preparation of ester derivatives of fatty acids for chromatographic analysis. *Advances in lipid methodology*, **2(69)**, e111.
- Clifton, P. M., and Keogh, J. B. (2017).** A systematic review of the effect of dietary saturated and polyunsaturated fat on heart disease. *Nutrition, Metabolism and Cardiovascular Diseases*, **27(12)**, 1060-1080.
- Díaz, G. C., Cruz, Y. R., Fortes, M. M., Viegas, C. V., Carliz, R. G., Furtado, N. C., and Aranda, D. A. G. (2014).** Primary Separation of Antioxidants (Unsaponifiables) the Wet Biomass Microalgae *Chlamydomonas* sp. and Production of the Biodiesel. *Natural Science*, **6(15)**, 1210.
- Dorni, C., Sharma, P., Saikia, G., and Longvah, T. (2018).** Fatty acid profile of edible oils and fats consumed in India. *Food chemistry*, **238**, 9-15.
- Eladel, H., Abomohra, A. E. F., Battah, M., Mohmmmed, S., Radwan, A., and Abdelrahim, H. (2019).** Evaluation of *Chlorella sorokiniana* isolated from local municipal wastewater for dual application in nutrient removal and biodiesel production. *Bioprocess and biosystems engineering*, **42(3)**, 425-433.
- Enamala, M. K., Enamala, S., Chavali, M., Donepudi, J., Yadavalli, R., Kolapalli, B., ... and Kuppam, C. (2018).** Production of biofuels from microalgae-A review on cultivation, harvesting, lipid extraction, and numerous applications of microalgae. *Renewable and Sustainable Energy Reviews*, **94**, 49-68.

- Folayan, A. J., Anawe, P. A. L., Aladejare, A. E., and Ayeni, A. O. (2019).** Experimental investigation of the effect of fatty acids configuration, chain length, branching and degree of unsaturation on biodiesel fuel properties obtained from lauric oils, high-oleic and high-linoleic vegetable oil biomass. *Energy Reports*, **5**, 793-806.
- Folch, J., Lees, M., and Stanley, G. S. (1957).** A simple method for the isolation and purification of total lipids from animal tissues. *Journal of biological chemistry*, **226(1)**, 497-509.
- Fraeye, I., Bruneel, C., Lemahieu, C., Buyse, J., Muylaert, K., and Foubert, I. (2012).** Dietary enrichment of eggs with omega-3 fatty acids: A review. *Food Research International*, **48(2)**, 961-969.
- Guihéneuf, F., and Stengel, D. B. (2013).** LC-PUFA-enriched oil production by microalgae: accumulation of lipid and triacylglycerols containing n-3 LC-PUFA is triggered by nitrogen limitation and inorganic carbon availability in the marine haptophyte *Pavlova lutheri*. *Marine drugs*, **11(11)**, 4246-4266.
- Hegewald, E., & Schmidt, A. (1992).** *Asterarcys Comas*, eine weit verbreitete tropische Grünalgenart. *Algological Studies/Archiv für Hydrobiologie, Supplement Volumes*, **25-30**.
- Held, P. (2011).** Monitoring of algal growth using their intrinsic properties: Use of a Multi-Mode Monochromator-based Microplate Reader for Biofuel Research.Inc., Vermont.

- Hishikawa, D., Valentine, W. J., Iizuka-Hishikawa, Y., Shindou, H., and Shimizu, T. (2017).** Metabolism and functions of docosahexaenoic acid-containing membrane glycerophospholipids. *FEBS letters*, **591(18)**, 2730-2744.
- Hoekman, S. K., Broch, A., Robbins, C., Cenicerros, E., and Natarajan, M. (2012).** Review of biodiesel composition, properties, and specifications. *Renewable and sustainable energy reviews*, **16(1)**, 143-169.
- Islam, M. A., Magnusson, M., Brown, R. J., Ayoko, G. A., Nabi, M., and Heimann, K. (2013).** Microalgal species selection for biodiesel production based on fuel properties derived from fatty acid profiles. *Energies*, **6(11)**, 5676-5702.
- Khan, M. I., Shin, J. H., & Kim, J. D. (2018).** The promising future of microalgae: Current status, challenges, and optimization of a sustainable and renewable industry for biofuels, feed, and other products. *Microbial Cell Factories*. <https://doi.org/10.1186/s12934-018-0879-x>
- Komárek, J., & Fott, B. (1983).** Die Binnengewässer, Band 16 Teil 7 Hälfte 1. Das Phytoplankton des Süßwassers. Systematik und Biologie Teil 7, 1. Hälfte: Chlorophyceae (Grünalgen), Ordnung Chlorococcales. *Schweizerbart'sche Verlagsbuchhandlung: Stuttgart, Germany*.
- Kramadibrata, M. A. M., Nurjanah, S., Muhaemin, M., Mardawati, E., Herwanto, T., Rosalinda, S. N., ... and Putri, F. E. (2019).** Selecting biofuel obtained from Sunan Pecan oil for diesel engine fuel. *Journal of Agricultural Science and Technology A*, **9**, 323-328.

- Liang, Y., Beardall, J., and Heraud, P. (2006).** Effects of nitrogen source and UV radiation on the growth, chlorophyll fluorescence and fatty acid composition of *Phaeodactylum tricornutum* and *Chaetoceros muelleri* (Bacillariophyceae). *Journal of Photochemistry and Photobiology B: Biology*, **82(3)**, 161-172.
- Ma, Y., Wang, Z., Yu, C., Yin, Y., & Zhou, G. (2014).** Evaluation of the potential of 9 *Nannochloropsis* strains for biodiesel production. *Bioresource technology*, **167**, 503-509.
- Maki, K. C., Eren, F., Cassens, M. E., Dicklin, M. R., and Davidson, M. H. (2018).**  $\omega$ -6 Polyunsaturated fatty acids and cardiometabolic health: current evidence, controversies, and research gaps. *Advances in Nutrition*, **9(6)**, 688-700.
- Mandotra, S. K., Kumar, P., Suseela, M. R., and Ramteke, P. W. (2014).** Fresh water green microalga *Scenedesmus abundans*: a potential feedstock for high quality biodiesel production. *Bioresource Technology*, **156**, 42-47.
- Mathimani, T., and Pugazhendhi, A. (2019).** Utilization of algae for biofuel, bio-products and bio-remediation. *Biocatalysis and agricultural biotechnology*, **17**, 326-330.
- Mendis, E., and Kim, S. K. (2011).** Present and future prospects of seaweeds in developing functional foods. *Advances in food and nutrition research*, **64**, 1-15.
- Meng, X., Yang, J., Xu, X., Zhang, L., Nie, Q., & Xian, M. (2009).** Biodiesel production from oleaginous microorganisms. *Renewable energy*, **34(1)**, 1-5.

- Park, J. Y., Kim, D. K., Lee, J. P., Park, S. C., Kim, Y. J., and Lee, J. S. (2008).** Blending effects of biodiesels on oxidation stability and low temperature flow properties. *Bioresource technology*, **99(5)**, 1196-1203.
- Ramos, M. J., Fernández, C. M., Casas, A., Rodríguez, L., and Pérez, Á. (2009).** Influence of fatty acid composition of raw materials on biodiesel properties. *Bioresource technology*, **100(1)**, 261-268.
- Santhakumaran, P., Kookal, S. K., Mathew, L., and Ray, J. G. (2019).** Bioprospecting of three rapid-growing freshwater green algae, promising biomass for biodiesel production. *BioEnergy Research*, **12(3)**, 680-693.
- Song, M., Pei, H., Hu, W., and Ma, G. (2013).** Evaluation of the potential of 10 microalgal strains for biodiesel production. *Bioresource technology*, **141**, 245-251.
- Talebi, A. F., Mohtashami, S. K., Tabatabaei, M., Tohidfar, M., Bagheri, A., Zeinalabedini, M., ... and Bakhtiari, S. (2013).** Fatty acids profiling: a selective criterion for screening microalgae strains for biodiesel production. *Algal Research*, **2(3)**, 258-267.
- Tiwari, V., Das, A., Thakur, S., & Trivedi, R. K. (2021).** Molecular characterization of blue-green algae (*Anabaena constricta*) and comparative studies of biodiesel production from other species.
- Tsarenko, P. M., Wasser, S. P., & Nevo, E. (2011).** Chlorophyta. *Algae of Ukraine: diversity, nomenclature, taxonomy, ecology and geography*. ARG Gantner, Ruggell, Liechtenstein.
- Ward, O. P., and Singh, A. (2005).** Omega-3/6 fatty acids: alternative sources of production. *Process biochemistry*, **40(12)**, 3627-3652.

- Wehr, J. D., Sheath, R. G., and Kociolek, J. P. (Eds.). (2015).** *Freshwater algae of North America: ecology and classification*. Elsevier.
- Wu, H., Xu, L., & Ballantyne, C. M. (2020).** Dietary and pharmacological fatty acids and cardiovascular health. *The Journal of Clinical Endocrinology & Metabolism*, **105(4)**, 1030-1045.
- Wysoczański T, Sokoła-Wysoczańska E, Pękala J, Lochyński S, Czyż K, Bodkowski R, Herbinger G, Patkowska-Sokoła B, Librowski T (2016).** Omega-3 fatty acids and their role in central nervous system a review. *Current medicinal chemistry*, **23(8)**, 816-831.
- Yagi, S., Fukuda, D., Aihara, K. I., Akaike, M., Shimabukuro, M., and Sata, M. (2017).** n-3 polyunsaturated fatty acids: promising nutrients for preventing cardiovascular disease. *Journal of atherosclerosis and thrombosis*, RV17013.
- Yen, H. W., Hu, I. C., Chen, C. Y., Ho, S. H., Lee, D. J., and Chang, J. S. (2013).** Microalgae-based biorefinery—from biofuels to natural products. *Bioresource technology*, **135**, 166-174.

## فحص إمكانية استخدام أربعة طحالب دقيقة خضراء كمواد أولية لإنتاج وقود الديزل الحيوي والمغذيات

سها محمد ، حامد العادل ، محمد بطاح ، ياسمين ابراهيم

قسم النبات والأحياء الدقيقة، كلية العلوم، جامعة بنها، بنها 13518، مصر

يُعد الكشف عن الطحالب الدقيقة الزيتية ذات الإنتاجية العالية للدهون ميزة أساسية لإنتاج الوقود الحيوي والمغذيات. لذلك فقد أصبح فحص واستكشاف الطحالب الدقيقة من البيئات المختلفة أمراً مهماً للغاية لمواجهة الطلب على هذه المصادر الحيوية. وبناءً عليه ، هدفت الدراسة الحالية إلى فحص كفاءة المواد الأولية للديزل الحيوي والقدرة على تجميع الأحماض الدهنية الأساسية لأربعة طحالب دقيقة خضراء هي: *Chlorolobion braunii* – *Tetrademus obliquus* - *Monoraphidium miutum* - *Asterarcys quadricellulare* ، والمعزولة من عينات التربة من بنها ، مصر. على الرغم من أن *T.obliquus* كان أعلى الأنواع المنتجة للكتلة الحيوية (114,86 مجم لتر / يوم) ، سجل *M. minutum* أعلى إنتاجية للدهون (29 مجم لتر / يوم). أكد محتوى الأحماض الدهنية للطحالب الأربعة ملاءمتها لإنتاج وقود الديزل الحيوي حيث سجلت خصائص وقود الديزل الحيوي ضمن حدود المعايير الدولية. أيضاً ، كما كشف اختبار الأنواع الأكثر ملائمة من خلال تحليل PROMETHEE و GAIA أن طحلب *C. braunii* هو الأفضل كمصدر محتمل لإنتاج وقود الديزل الحيوي لأنه يحتوي على نسبة عالية من الأحماض الدهنية المشبعة (39.75%) ويظهر رقم السيتان (54.99) وقيمة اليود المنخفضة (100.57) والعديد من خصائص وقود الديزل الحيوي الأخرى ضمن حدود المعايير الدولية. بالإضافة إلى ذلك ، أظهرت الطحالب الدقيقة التي تم فحصها نسبة عالية من أحماض الأوميغا الدهنية في الأحماض الدهنية الكلية. حيث سجل كلا من طحلب *Asterarcys quadricellulare* و *M. minutum* نسبة عالية من الأحماض الدهنية أوميغا 3 بنسبة 34.4% و 28.9% على التوالي.